

Imperata cylindrica may be useful for soil erosion control due to the fast growth of its rhizomes. It may also have some potential in silvi-pastoralist systems as it can grow in a relatively low light. For example it is a suitable grass in coconut plantations ([Senanayake, 1995](#)).

Datasheet citation

Heuzé V., Tran G., Baumont R., Bastianelli D., 2016. *Alang-alang (Imperata cylindrica)*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/425> Last updated on March 17, 2016, 9:26

English correction by Tim Smith (Animal Science consultant) and Hélène Thiolllet (AFZ)


Image credits

● Gilles Tran, AFZ ● Gilles Tran, AFZ ● Gilles Tran, AFZ

 Share / Save   



Automatic translation

 Sélectionner une langue ▼

Feed categories

- All feeds
- Forage plants
 - ▶ Cereal and grass forages
 - ▶ Legume forages
 - ▶ Forage trees
 - ▶ Aquatic plants
 - ▶ Other forage plants
- Plant products/by-products
 - ▶ Cereal grains and by-products
 - ▶ Legume seeds and by-products
 - ▶ Oil plants and by-products
 - ▶ Fruits and by-products
 - ▶ Roots, tubers and by-products
 - ▶ Sugar processing by-products
 - ▶ Plant oils and fats
 - ▶ Other plant by-products
- Feeds of animal origin
 - ▶ Animal by-products
 - ▶ Dairy products/by-products
 - ▶ Animal fats and oils
 - ▶ Insects
- Other feeds
 - ▶ Minerals
 - ▶ Other products

Latin names

- Plant and animal families
- Plant and animal species

Resources

- Broadening horizons
- Literature search
- Image search
- Glossary
- External resources
 - ▶ Literature databases
 - ▶ Feeds and plants databases
 - ▶ Organisations & networks
 - ▶ Books
 - ▶ Journals

Alang-alang (Imperata cylindrica)

- Description
- Nutritional aspects
- Nutritional tables
- References

Nutritional attributes

Imperata cylindrica is a poor quality grass. Its protein content is about 7% DM and generally lower than 12% DM (Feedipedia, 2013). It is also deficient in energy and sodium (Falvey, 1981).

Potential constraints

Cattle tend to avoid mature leaves due to their sharp tips and edges that cause mouth irritation (Soerjani, 1970).

Ruminants

Though *Imperata cylindrica* pastures may be used profitably in some situations, they must be supplemented with other pasture species and feeds to obtain acceptable animal performance (Falvey, 1981; Holmes et al., 1980). In the Thai Highlands, *Imperata cylindrica* provides some forage but tends to decline or disappear if continually grazed by cattle (Andrews, 1983).

Nutritive value

It has been estimated that the low protein content of *Imperata cylindrica* forage can support live weight gain in beef cattle only for the first 6 weeks of forage growth (Papua New Guinea) or 20 weeks (Thai Highlands) (Falvey, 1981). Nylon bag digestibility of *Imperata cylindrica* was found to be 2/3 lower than that of buffel grass (*Cenchrus ciliaris*), *Setaria sphacelata* and elephant grass (*Pennisetum purpureum*) (Holmes et al., 1980). In the Thai Highlands, *in vitro* digestibility declined with plant age, from 70% when young to 40% at 150 days, and depended on the season: 50-70% for wet season regrowth and lower than 45% in the dry season (Falvey, 1981).

Palatability

Imperata cylindrica is palatable if cut frequently but the mature leaves are sharp and irritating (see Potential constraints above) (Soerjani, 1970).

Beef cattle

Imperata cylindrica pastures can support a viable extensive cattle production system provided that they are supplemented with sources of energy (carbohydrates such as cassava), nitrogen (urea or legumes), and minerals (Falvey, 1981; Holmes et al., 1980; Soewardi et al., 1974). Weight gains of cattle grazing non-supplemented *Imperata cylindrica* were reported to be lower than those obtained with supplemented *Imperata cylindrica* pasture, or with other pasture species at higher stocking rates.

Using urea-molasses-mineral blocks and introducing new fodder species (especially legumes) significantly improved production from *Imperata cylindrica* grasslands in smallholder farming systems. Fodder species for fallow improvement, modified alley cropping or hedgerow systems and plantations integrating livestock production may help to increase sustainability (Calub et al., 1996).

The following table presents several trials involving *Imperata cylindrica* pastures.

Animal	Region	Average daily gain and stocking rate for <i>Imperata cylindrica</i>	Average daily gain and stocking rate for other forage species or supplemented <i>Imperata cylindrica</i>	Reference
Heifers	Papua New Guinea	0.2-0.25 kg/day, 0.8-1.6 animal/ha	0.45 kg/day, 1.7-2.2 animals/ha on Guinea grass and legume pastures	Holmes et al., 1980
Steers	Papua New Guinea	0.38 kg/day	0.47-0.52-0.63 kg/day when supplemented with palm kernel meal alone, with molasses or with molasses and urea 0.53-0.56-0.54 kg/day when supplemented with palm kernel meal alone, with molasses or with molasses and urea	Gaigal et al., 2000
Cattle	Philippines	0.27 kg/day, 1 animal/ha	>0.81 kg/day, 2 animal/ha on <i>Brachiaria mutica</i> /centro pasture	Magadan et al., 1974
Cattle	Thai Highlands	0.04 kg/day	+30% with sodium supplementation	Falvey, 1981
Cattle	Thai Highlands	0.04 kg/day	0.21 kg/day with legume supplementation, 0.24 kg/day with urea-molasses and mineral block	Mikled, 1976
Cattle	Indonesia	0.17 kg/day	0.21 kg/day with urea, carbohydrate and mineral supplementation	Soewardi et al., 1974




Sheep and goats

Supplementing *Imperata cylindrica* with legumes is a valuable strategy in small ruminants. Dry matter intakes of 2.8 and 3.3 % LW (for goats and sheep respectively) were obtained using alang-alang grass (*ad libitum*) and *Leucaena leucocephala* (300 g). These values were higher than those obtained for *I. cylindrica* or *Leucaena* alone. Diet digestibility of the low-quality herbage was higher for goats than for sheep (Komolong et al., 1988).

Pigs

Early researchers noted that the succulent white and starchy rhizomes were eaten and even actively searched for by pigs (Hole, 1911). However, later attempts to fatten pigs on the rhizomes have failed as the pigs lost weight (Hubbard, 1944).

Poultry



The composition of *Imperata cylindrica* makes it unsuitable as feed ingredient for poultry. However, in an experiment on young chickens, supplementation of small quantities (2 to 6% DM) of *Imperata cylindrica* to the basal diet resulted in improved performance ([Kencana et al., 1980](#)).

Rabbits

No information found (2013).

Datasheet citation

Heuzé V., Tran G., Baumont R., Bastianelli D., 2016. *Alang-alang (Imperata cylindrica)*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/425> Last updated on March 17, 2016, 9:26

Image credits

- Gilles Tran, AFZ ● Gilles Tran, AFZ ● Gilles Tran, AFZ

Share / Save



Alang-alang (Imperata cylindrica)

- Description
- Nutritional aspects
- Nutritional tables
- References

Tables of chemical composition and nutritional value

- Alang-alang (Imperata cylindrica), aerial part, fresh
- Alang-alang (Imperata cylindrica), hay

Avg: average or predicted value; SD: standard deviation; Min: minimum value; Max: maximum value; Nb: number of values (samples) used

Alang-alang (Imperata cylindrica), aerial part, fresh



Main analysis	Unit	Avg	SD	Min	Max	Nb
Dry matter	% as fed	31.9	5.0	23.8	39.2	16
Crude protein	% DM	6.5	2.0	3.5	11.2	21
Crude fibre	% DM	39.4	4.1	32.1	44.9	21
NDF	% DM	74.3	2.8	70.1	77.2	7 *
ADF	% DM	45.7	1.8	41.5	46.8	7 *
Lignin	% DM	6.6	0.9	6.6	10.2	7 *
Ether extract	% DM	1.9	0.5	1.3	3.0	21
Ash	% DM	7.0	1.4	4.7	9.0	21
Gross energy	MJ/kg DM	18.6				*

Minerals	Unit	Avg	SD	Min	Max	Nb
Calcium	g/kg DM	3.3	3.5	0.9	17.4	19
Phosphorus	g/kg DM	1.4	0.6	0.4	2.6	19
Potassium	g/kg DM	11.7	5.3	3.6	19.1	15
Sodium	g/kg DM	0.2	0.2	0.1	0.5	3
Magnesium	g/kg DM	2.1	1.0	0.9	4.2	13
Manganese	mg/kg DM	89	79	18	193	4
Zinc	mg/kg DM	13	4	9	19	4
Copper	mg/kg DM	3	1	3	4	4

Ruminant nutritive values	Unit	Avg	SD	Min	Max	Nb
OM digestibility, Ruminant	%	57.2				*
Energy digestibility, ruminants	%	54.7				*
DE ruminants	MJ/kg DM	10.1				*
ME ruminants	MJ/kg DM	8.2				*
Nitrogen digestibility, ruminants	%	30.0				1

The asterisk * indicates that the average value was obtained by an equation.

References

CIRAD, 1991; Laksevela et al., 1970; Lim Han Kuo, 1967; Nasrullah et al., 2003; Pozy et al., 1996; Senanayake, 1995

Last updated on 24/10/2012 00:43:34

Alang-alang (Imperata cylindrica), hay



Main analysis	Unit	Avg	SD	Min	Max	Nb
Crude protein	% DM	3.8				1
Crude fibre	% DM	39.7				1
NDF	% DM	74.6				*
ADF	% DM	46.0				*
Lignin	% DM	6.7				*
Ether extract	% DM	0.7				1

Ash	% DM	7.8	1			
Gross energy	MJ/kg DM	18.0	*			
Ruminant nutritive values	Unit	Avg	SD	Min	Max	Nb
OM digestibility, Ruminant	%	52.9				*
Energy digestibility, ruminants	%	49.5				*
DE ruminants	MJ/kg DM	8.9				*
ME ruminants	MJ/kg DM	7.2				*
Nitrogen digestibility, ruminants	%	34.0				1

The asterisk * indicates that the average value was obtained by an equation.

References

Sen, 1938

Last updated on 24/10/2012 00:43:34

Datasheet citation

Heuzé V., Tran G., Baumont R., Bastianelli D., 2016. *Alang-alang (Imperata cylindrica)*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/425>. Last updated on March 17, 2016, 9:26

English correction by Tim Smith (Animal Science consultant) and Hélène Thiolllet (AFZ)

Image credits

- Gilles Tran, AFZ
- Gilles Tran, AFZ
- Gilles Tran, AFZ



Alang-alang (Imperata cylindrica)

- Description
- Nutritional aspects
- Nutritional tables
- References

References

Andrews, A. C., 1983. *Imperata cylindrica* in the highlands of northern Thailand: its productivity and status as a weed. Mountain Res. Devel., 3 (4): 386-388

Calub, A. D. ; Anwarhan, H. ; Roder, W., 1996. Livestock production systems for Imperata grasslands. Agroforestry Systems, 36 (1-3): 121-128

Chadhokar, P.A., 1977. Establishment of stylo (*Stylosanthes guianensis*) in kunai (*Imperata cylindrica*) pastures and its effects on dry matter yield and animal production in the Markham valley, Papua New Guinea. Trop. Grassl. 11 (3):263-272

Ecocrop, 2010. Ecocrop database. FAO

Ecoport, 2010. Ecoport database. Ecoport

Falvey, L.: Andrew, A., 1979. Improved pastures in the Thai highlands. Trop. Grassl., 13: 154-156

Falvey, L., 1977. Review of existing knowledge of ruminants in the highlands of Northern Thailand. Thai J. Agric. Sci., 10: 111-119

Falvey, J. L., 1981. *Imperata cylindrica* for animal production in South-East Asia: a review. Trop. Grassl., 15 (1): 52-56

FAO, 2010. Grassland Index. A searchable catalogue of grass and forage legumes. FAO

Friday, K. S. ; Drilling, M. E. ; Garrity, D. P., 1999. *Imperata* grassland rehabilitation using agroforestry and assisted natural regeneration. International Centre for Research in Agroforestry, Southeast Asian Regional Research Programme Bogor, Indonesia 1999

Galgai, K. K. ; Komolung, M. K., 2000. Copra meal and palm kernel meal supplementation with and without molasses and urea to weaner steers grazing *Imperata cylindrica* pastures in Papua New Guinea. Asian-Aust. J. Anim. Sci., 13 (Suppl. July 2000): 261

Gbehounou, G. ; Ouédraogo, O. ; Tehia, E. ; Chikoye, D., 2000. Rapport des pays: Bénin, Burkina-Faso, Côte d'Ivoire, Nigéria. in: Atelier de lutte contre l'Imperata cylindrica, Labrada, R. ed. Direction de la Production et Protection Végétale de la FAO (AGP), Rome

GISD, 2010. Global Invasive Species Database. Invasive Species Specialist Group of the IUCN

Hole, R. S., 1911. On some Indian forest grasses and their ecology. Indian Forest Memoirs. Bot. Ser. Part I, 1-26

Holmes, J. H. G. ; Lemerle, C. ; Schottler, J. H., 1980. *Imperata cylindrica* for cattle production in Papua New Guinea. Papua New Guinea Agric. J., 31 (1/4): 51-62

Hubbard, C. E., 1944. *Imperata cylindrica*: Taxonomy, distribution, economic significance and control. Imperial Agricultural Bureaux Joint Publication No. 7. Oxford, UK: Imperial Forestry Bureau; Aberystwyth, UK: Imperial Bureau of Pastures and Forage Crops. 63 p.

Jonathan, J. ; Hariadi, B.P.J., 1999. *Imperata cylindrica* (L.) Raeuschel. Record from Proseabase. de Padua, L.S., Bunyapraphatsara, N. and Lemmens, R.H.M.J. (Editors). PROSEA (Plant Resources of South-East Asia) Foundation, Bogor, Indonesia.

Kencana, S. ; Hartiko, H., 1980. The use of alang-alang (*Imperata cylindrica* (L.) Beauv.) as green supplement in broiler chicken feed. (A preliminary study). Proceedings of BIOTROP Workshop on Alang-alang, Bogor (1976). 1980. 231-236

Komolung, M. ; Sriskandarajah, N., 1988. The use of *Imperata cylindrica* (L.) Beauv. by grazing cattle in Papua-New Guinea. Proceedings of Papua New Guinea Society of Animal Production, Lae Morobe Province, June 1988. Maximising Animal Production in Papua New Guinea.

Labrada, R., 2003. Weed Management for Developing Countries. Addendum 1. Plant production and protection papers N °120. FAO, Rome

Laksevela, B. ; Said A. N., 1970. Kenya sisal project. Bull. No. 71, Kenya Sisal Board, p. 13

Lim Han Kuo, 1967. Animal feeding stuffs. Part 3. Compositional data of feeds and concentrates. Malay. Agric. J., 46 (1): 63-79

Macdicken, K. G. ; Hairiah, K. ; Otsamo, A. ; Duguma, B. ; Majid, N.M., 1996. Shade-based control of *Imperata cylindrica*: tree fallows and cover crops. Agroforestry systems, 36 (1-3): 131-149

Magadan, P. B. ; Javier, E. Q. ;Madamba, J. C., 1974. Beef production on native (*Imperata cylindrica* (L.) Beauv.) and para grass (*Brachiaria mutica* (Forsk.) Stapf) pastures in the Philippines. Proc. 12th Int. Grassl. Congr., Vol. 3, part 1, 293-298

Mikled, C., 1976. Dry season supplementation of cattle grazing native grasslands in the North Thailand highlands. Thai J. Agric. Sci., 10 (3): 135-142

Moog, F. A., 1983. Beef production on Leucaena Imperata pastures and cattle feeds on small farms in the Philippines. Leucaena research in the Asian Pacific region. 1983, 69-72. Ottawa, Ontario, Canada

Peet, N. B. ; Watkinson, A. R. ; Bell, D. J. ; Sharma, U. R., 1999. The conservation management of *Imperata cylindrica* grassland in Nepal with fire and cutting: an experimental approach. J. Appl. Ecol., 36 (3): 374-387

Pomplona, P., 1974. Alang alang can be processed to pellets for animal feed. BioTrop Newsletter, 9: 15-16

Rahman, M. ; Subagyo, T., 2000. The effect of alang-alang (*Imperata cylindrica* (L.) Beauv.) leaves on some upland crops and dead materials of kiambang (*Salvinia molesta* D.S. Mitchell) on lowland rice seedlings. Proc. 3rd Indon. Weed

Science Conf., Bandung

Sajise, P.E. ; Lales, J.S., 1975. Allelopathy in a mixture of cogon (*Imperata cylindrica*) and *Stylosanthes guyanensis*. *Kalikasan, Philippine J. Biol.*, 4 (2): 155-164

Sen, K. C., 1938. The nutritive values of Indian cattle feeds and the feeding of animals. Indian Council of Agricultural Research, New Dehli, Bulletin No. 25, 1-30

Senanayake, S. G. J. N., 1995. The effects of different light levels on the nutritive quality of four natural tropical grasses. *Trop. Grassl.*, 29 (2): 111-114

Soerjani, M., 1970. Alang-alang. Pattern of growth as related to the problem of control. Bogor, Indonesia. SEAMEO Regional Center for Tropical Biology. Biotrop. Bull. No. 1.

Soewardi, B. ; Sastradipradja, D. ; Nosoetion, A. H. ; Hutasoit, J. H., 1974. Studies on alang-alang (*Imperata cylindrica* (L.) Beauv.) for cattle feeding. 1. The effects of carbohydrate sources in urea containing concentrate on feed utilization. *Biotrop Bulletin*, 8: 25

USDA, 2010. GRIN - Germplasm Resources Information Network. National Germplasm Resources Laboratory, Beltsville, Maryland

USDA, 2013. GRIN - Germplasm Resources Information Network. National Germplasm Resources Laboratory, Beltsville, Maryland

36 references found

Datasheet citation

Heuzé V., Tran G., Baumont R., Bastianelli D., 2016. *Alang-alang (Imperata cylindrica)*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/425>. Last updated on March 17, 2016, 9:26

English correction by Tim Smith (Animal Science consultant) and Hélène Thiollet (AFZ)

Image credits

- Gilles Tran, AFZ
- Gilles Tran, AFZ
- Gilles Tran, AFZ

